


AOP Development; Aromatase Inhibition and Androgen Receptor Agonism Lead to Male Biased Sex Ratio Via Impacts on Gonad Differentiation

- potential adverse effects of concern.
- ecological risk assessment.

- with environmental sex differentiation (ESD)



K.J. Santana Rodriguez¹, D. L. Villeneuve², G. T. Ankley², D. H. Miller² ¹Oak Ridge Institute for Science and Education (ORISE), ²U.S. EPA Great Lakes Toxicology and Ecology Division

Contents of this poster neither constitute nor necessarily reflect US EPA policy.



For reference list please email santanarodriguez.kelvin@epa.gov

- **上**. 2. https://doi.org/10.1016/j.aquatox.2016.04.018 3. Tilapia. Endocrinology. 158. 10.1210/en.2017-00127. 4. 5. 6. 6433(03)00178-8 7. 8. 9. Miller et al 2021 10. 11. 12. https://doi.org/10.1210/en.2017-00617 13. 14. 15. 16. 17.

REFERENCES

Lau, E. S., Zhang, Z., Qin, M., & Ge, W. (2016). Knockout of Zebrafish Ovarian Aromatase Gene (cyp19a1a) by TALEN and CRISPR/Cas9 Leads to All-male Offspring Due to Failed Ovarian Differentiation. Scientific reports, 6, 37357. https://doi.org/10.1038/srep37357

Muth-Köhne, E., Westphal-Settele, K., Brückner, J., Konradi, S., Schiller, V., Schäfers, C., Teigeler, M. (2016). Linking the response of endocrine regulated genes to adverse effects on sex differentiation improves comprehension of aromatase inhibition in a Fish Sexual Development Test. Aquatic toxicology (Amsterdam, Netherlands), 176, 116–127.

Zhang, Xianbo & Li, Mengru & Ma, He & Liu, Xingyong & Shi, Hongjuan & Li, Minghui & Wang, Deshou. (2017). Mutation of foxl2 or cyp19a1a Results in Female to Male Sex Reversal in XX Nile

Holbech, H., Kinnberg, K. L., Brande-Lavridsen, N., Bjerregaard, P., Petersen, G. I., Norrgren, L., Örn, S., Braunbeck, T., Baumann, L., Bomke, C., Dorgerloh, M., Bruns, E., Ruehl-Fehlert, C., Green, J. W., Springer, T. A., & Gourmelon, A. (2012). Comparison of zebrafish (Danio rerio) and fathead minnow (Pimephales promelas) as test species in the Fish Sexual Development Test (FSDT). Comparative Biochemistry and Physiology - C Toxicology and Pharmacology, 155(2), 407–415. https://doi.org/10.1016/j.cbpc.2011.11.002 Kwon, J. Y., Haghpanah, V., Kogson-Hurtado, L. M., McAndrew, B. J., & Penman, D. J. (2000). Masculinization of genetic female nile tilapia (Oreochromis niloticus) by dietary administration of an aromatase inhibitor during sexual differentiation. The Journal of experimental zoology, 287(1), 46–53. Uchida, D., Yamashita, M., Kitano, T., & Iguchi, T. (2004). An aromatase inhibitor or high water temperature induce oocyte apoptosis and depletion of P450 aromatase activity in the gonads of genetic female zebrafish during sex-reversal. Comparative biochemistry and physiology. Part A, Molecular & integrative physiology, 137(1), 11–20. https://doi.org/10.1016/s1095-

Ruksana, S., Pandit, N. P., & Nakamura, M. (2010). Efficacy of exemestane, a new generation in a gonochoristic fish. Comparative biochemistry and physiology. Toxicology & pharmacology : CBP, 152(1), 69–74. https://doi.org/10.1016/j.cbpc.2010.02.014 Thorpe, K. L., Marca Pereira, M. L., Schiffer, H., Burkhardt-Holm, P., Weber, K., & Wheeler, J. R. (2011). Mode of sexual differentiation and its influence on the relative sensitivity of the fathead minnow and zebrafish in the fish sexual development test. Aquatic Toxicology, 105(3–4), 412–420. https://doi.org/10.1016/j.aquatox.2011.07.012 Yin, Y., Tang, H., Liu, Y., Chen, Y., Li, G., Liu, X., & Lin, H. (2017). Targeted Disruption of Aromatase Reveals Dual Functions of cyp19a1a During Sex Differentiation in

Zebrafish. *Endocrinology*, 158(9), 3030–3041. https://doi.org/10.1210/en.2016-1865

Brown, A. R., Owen, S. F., Peters, J., Zhang, Y., Soffker, M., Paull, G. C., Hosken, D. J., Wahab, M. A., & Tyler, C. R. (2015). Climate change and pollution speed declines in zebrafish populations. Proceedings of the National Academy of Sciences of the United States of America, 112(11), E1237–E1246. https://doi.org/10.1073/pnas.1416269112 Crowder, C. M., Lassiter, C. S., & Gorelick, D. A. (2018). Nuclear Androgen Receptor Regulates Testes Organization and Oocyte Maturation in Zebrafish. Endocrinology, 159(2), 980–993.

Yu, G., Zhang, D., Liu, W., Wang, J., Liu, X., Zhou, C., Gui, J., & Xiao, W. (2018). Zebrafish androgen receptor is required for spermatogenesis and maintenance of ovarian function. Oncotarget, 9(36), 24320–24334. https://doi.org/10.18632/oncotarget.24407

Orn, S., Holbech, H., Madsen, T. H., Norrgren, L., & Petersen, G. I. (2003). Gonad development and vitellogenin production in zebrafish (Danio rerio) exposed to ethinylestradiol and methyltestosterone. Aquatic toxicology (Amsterdam, Netherlands), 65(4), 397–411. https://doi.org/10.1016/s0166-445x(03)00177-2 Morthorst, J. E., Holbech, H., & Bjerregaard, P. (2010). Trenbolone causes irreversible masculinization of zebrafish at environmentally relevant concentrations. Aquatic toxicology (Amsterdam, Netherlands), 98(4), 336–343. https://doi.org/10.1016/j.aquatox.2010.03.008

Baumann, L., Holbech, H., Keiter, S., Kinnberg, K. L., Knörr, S., Nagel, T., & Braunbeck, T. (2013). The maturity index as a tool to facilitate the interpretation of changes in vitellogenin production and sex ratio in the Fish Sexual Development Test. Aquatic toxicology (Amsterdam, Netherlands), 128-129, 34–42. Baumann, L., Knörr, S., Keiter, S., Nagel, T., Rehberger, K., Volz, S., Oberrauch, S., Schiller, V., Fenske, M., Holbech, H., & Braunbeck, T. (2014). Persistence of endocrine disruption in zebrafish (Danio rerio) after discontinued exposure to the androgen 17β-trenbolone. Environmental toxicology and chemistry, 33(11), 2488–2496. https://doi.org/10.1002/etc.2698