



WWW.EPA.GOV

# High-Throughput Assessment of Increased Chemical Toxicity Due to Hepatic Steatosis

Nyssa N. Tucker<sup>1,2</sup>, Gail M. Nelson<sup>2</sup>, Joshua A. Harrill<sup>2</sup>, Brian N. Chorley<sup>2\*</sup>

<sup>1</sup>Oak Ridge Institute for Science and Education, Oak Ridge, TN 37831

<sup>2</sup>Center for Computational Toxicology & Exposure, US Environmental Protection Agency, RTP, NC 27709

P 130

# 2137



\*Brian N. Chorley | Chorley.Brian@epa.gov | 919-541-2329

## Abstract

Hepatic steatosis alters native liver xenobiotic metabolism, impacts the bioactivation or detoxification of chemicals, and may alter sensitivity to chemical toxicity. Here, we modeled this state in a human hepatic cell culture model to assess the impact of steatosis on chemical toxicity in a quantitative and high-throughput manner. We induced steatosis in human hepatoma-derived cells, HepaRGs, by dosing maintenance media with 1 mM 1:2 oleic:palmitic free fatty acid for 1 week. Cytochrome P450 (CYP) gene expression and metabolic activity (CYPs 1A1, 1A2, 2B6, 2C9, 2E1, 3A4) was significantly altered in the steatotic culture condition. Relative culture viability was determined by CellTiter-Glo (CTG), lactate dehydrogenase (LDH) release, and multiplexed fluorometric measurements of nuclear morphology using the Opera Phenix high-content screening (HCS) system. Naive and steatotic HepaRG cells were exposed to known hepatotoxicant (rotenone) over a 5-point dose range for 24 or 48 hrs. Rotenone toxicity (IC50) shifted from a baseline 0.64  $\mu$ M to 0.48  $\mu$ M in steatotic cells as measured by CTG, from 0.83  $\mu$ M to 0.57  $\mu$ M measuring LDH, and 0.80  $\mu$ M to 0.62  $\mu$ M using cell counts derived from HCS. Additional toxicity measures - including morphology, reactive oxygen species generation, and mitochondrial membrane potential - as well as additional chemicals known to be impacted by CYP-mediated metabolism are currently being assessed. The results of this study indicated that we can measure the impact of pre-existing conditions on environmental chemical toxicity, which will contribute to ongoing Agency efforts to assess environmental exposure risks in susceptible subpopulations.

*This poster does not necessarily reflect EPA policy. Mention of trade names is not an endorsement or recommendation for use.*



## Introduction

Hepatic Steatosis = Fatty liver

○ Prevalent ~1/3<sup>rd</sup> of the world

○ Multiple causes →

A) Genetic B) Behavioral C) Exposure




○ Often asymptomatic & reversible

○ Causes morphological cellular changes


○ Disease state (altered lipid metabolism +  $\uparrow$  triglyceride retention) may increase susceptibility to environmental chemicals.

## In Vitro Model




**Figure 1. Establishing significant fat build-up in HepaRG cells.** Oil red O staining indicated intracellular accumulation of lipid droplets after free-fatty acid [FFA] exposure in the media for 1 week. The addition of a known hepatotoxicant and mitochondrial respiratory chain complex I inhibitor, significantly increased fat retention due to decreased fatty acid metabolism.

## R1: Model Characterization




**Figure 2. Fluorometric indication of hepatic steatosis in HepaRG cells.** Hoechst [HO] nuclear stain and Nile Red [NR] triglyceride stain following 48h exposure to media containing 1 mM of a 1:2 oleate:palmitate free-fatty acid [FFA].


## R2: Altered Susceptibility + Toxic Fingerprint




**Figure 6. Cellular membrane potential change with low-concentration test chemical exposure.** HepaRG with no FFA exposure, 48h 0.2  $\mu$ M rotenone with a HO nuclear stain and TMRM (mitochondrial membrane potential) / TOTO-3 (cellular membrane integrity).



**Figure 3. Quantitative lipid accumulation with NR staining.** Lipid accumulation due to 1 week of FFA exposure in media was assessed by measuring fluorescence of lipid-bound Nile Red dye normalized to cell count determined by Hoechst staining.




**Figure 4. CYP gene expression changes due to hepatic steatosis in HepaRG.** Real-time qPCR measured 6 common CYP enzymes active in human liver. Metabolic enzyme expression shown as fold-change following 7-day incubation of 1 mM 1:2 oleate:palmitate FFA.




**Figure 8. Comparative HT measures with 24 h test chemical exposure over dose range.** Using the Opera Phenix imager and Harmony software, fluorescent stains are reduced to values representing single cell data resolution. Further analysis using R, R studio, and the TidyVerse suite result in quantitative outputs confirming visual and complementary assay outcomes.

## R3: Correlation of Low- and High- Throughput Measures



**Figure 10. Rotenone-mediated toxicity increases when cells are steatotic.** Data points derived from an ATP-proxy viability assay were fit to a non-linear least-squares model and IC50 values were determined ([www.ic50.tk](http://www.ic50.tk); see Figure 10).

## WIP: Interrogating CYPs with Targeted Chemicals



**Figure 12. Nuclear morphological assessments with 24 h high concentration test chemical exposure.** Two methods of FFA exposure with test chemical produce visually distinct outcomes, potentially due to differences in chemical uptake and processing. Examining nuclear morphology indicates potential use of fluorescent intensity as proxy measures of cell viability. Further analysis of textural features are potentially useful indicators of sub-toxic responses.

## Conclusion and Future Directions

- Steatotic HepaRG: viable but have altered CYP metabolism.
- This state **alters the hepatotoxicity** of some chemical exposures, underscoring the importance of assessing hepatic steatosis as a **common susceptibility factor** for chemical toxicity.
- We can assess the impact of **CYP enzyme alteration due to steatosis** by examining reference toxicants targeted by specific CYP metabolism:
  - High-content imaging/analysis, especially **GIS topographical measures**, will help us assess the impact of hepatic steatosis on chemical-mediated toxicity;
- Mitochondrial or cellular membrane potentials coupled with GSH/ROS measures may indicate **additional parameters of mechanistic toxicity** using similar workflow processes.

Greatest thanks to Denise MacMillan and the chemistry core at the EPA for HPLC ID of CYP activity! Same to my collaborators / mentor! Literally couldn't do it alone!

U.S. Environmental Protection Agency  
Office of Research and Development  
Center for Computational Toxicology & Exposure  
Biomolecular & Computational Toxicology Division  
Alternative Experimental Toxicology Models Branch

Fatty liver changes liver metabolism.  
The resultant impact on chemical toxicity can be quantified in a high-throughput, hepatic cell culture.